



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

CYCLOTOMIC HEPTASECTION FOR THE PRIME 43.

BY PANDIT OUDH UPADHYAYA.*

The problem of cyclotomic section has engaged the attention of many eminent mathematicians and solutions have been obtained by them for particular cases. The problem of the trisection and quartisection was completely solved by Cayley in a paper in which he also discussed the quinquisection but did not complete the solution. He once again took up the same problem in the proceedings of the London Mathematical Society in 1881 but was not able to complete the solution.

The problem of quinquisection was first solved by Rogers.† The same problem has very recently been solved by Burnside.‡ Towards the end of his paper he refers to the case of heptasection and says "I have carried the case $q = 7$ so far as to assure myself that it is not quite parallel with that of $q = 5$; a set of three simultaneous Diophantine equations occur, but they are not sufficient to ensure that the equations expressing the product of A 's form a consistent multiplication table." In view of this statement it is believed that the problem of heptasection for the prime 43 has not been previously considered.

The object of this paper is to consider the problem of heptasection for the prime 43. All the details of calculation have been suppressed in order to save space, and only the final result is given.

Let a be an imaginary root of $x^{43} - 1 = 0$, and let us divide all the imaginary roots into 7 groups according to the following scheme:

$$\begin{aligned} A &= a + a^{42} + a^{37} + a^6 + a^{36} + a^7, \\ B &= a^3 + a^{40} + a^{25} + a^{18} + a^{22} + a^{21}, \\ C &= a^9 + a^{34} + a^{32} + a^{11} + a^{23} + a^{20}, \\ D &= a^{27} + a^{16} + a^{10} + a^{33} + a^{26} + a^{17}, \\ E &= a^{38} + a^5 + a^{30} + a^{13} + a^{35} + a^8, \\ F &= a^{28} + a^{15} + a^4 + a^{39} + a^{19} + a^{24}, \\ G &= a^{41} + a^2 + a^{12} + a^{31} + a^{14} + a^{29}. \end{aligned}$$

Calculating the elementary symmetric functions of these expressions we get:

$$\begin{aligned} \Sigma A &= -1, & \Sigma AB &= -18, & \Sigma ABC &= 35, & \Sigma ABCD &= 38, \\ \Sigma ABCDE &= -104, & \Sigma ABCDEF &= 7, & ABCDEFG &= 49. \end{aligned}$$

* Babu Shiva Prasad Gupta Research Scholar.

† Lond. Math. Soc. Proc., vol. 32 (1900-01), pp. 199-207.

‡ Lond. Math. Soc. Proc., vol. (2) 14, (1915), pp. 251-259.

The equation whose roots are A, B, C, D, E, F, G is therefore

$$\eta^7 + \eta^6 - 18\eta^5 - 35\eta^4 + 38\eta^3 + 104\eta^2 + 7\eta - 49 = 0.$$

Every root of this equation may be expressed as a rational integral function of any one assigned root; it is therefore an Abelian equation and can be solved by radicals.

I should like to mention that I have received a great amount of help in calculation from Pandit Shukdeo Chaube, Babu Brahmdeo Roy, Babu Raichand Bothera, and Sohan Lal Dugar.